arXiv:math/0702171v1 [math.AP] 7 Feb 2007

REMOVABLE SINGULARITY OF THE POLYHARMONIC EQUATION

Shu-Yu Hsu

Department of Mathematics
National Chung Cheng University
168 University Road, Min-Hsiung
Chia-Yi 621, Taiwan, R.O.C.
e-mail:syhsu@math.ccu.edu.tw

Feb 7, 2007

Abstract. Let $x_0 \in \Omega \subset \mathbb{R}^n$, $n \ge 2$, be a domain and let $m \ge 2$. We will prove that a solution u of the polyharmonic equation $\Delta^m u = 0$ in $\Omega \setminus \{x_0\}$ has a removable singularity at x_0 if and only if $|\Delta^k u(x)| = o(|x-x_0|^{2-n}) \quad \forall k = 0,1,2,\dots,m-1$ as $|x-x_0| \to 0$ for $n \ge 3$ and $= o(\log(|x-x_0|^{-1})) \quad \forall k = 0,1,2,\dots,m-1$ as $|x-x_0| \to 0$ for $n = 2$. For $m \ge 2$ we will also prove that u has a removable singularity at x_0 if $|u(x)| = o(|x-x_0|^{2m-n})$ as $|x-x_0| \to 0$ for $n \ge 3$ and $|u(x)| = o(|x-x_0|^{2m-2}\log(|x-x_0|^{-1}))$ as $|x-x_0| \to 0$ for $n = 2$.

Recently there is a lot of study of the singularities of equations. Singularities of solutions of the Ricci flow was studied by R.S. Hamilton [H] and G. Perelman [P1],[P2]. Removable singularities of the solution of the Ricci flow equation on $\mathbb{R}^2$ was studied by S.Y. Hsu in [Hs1]. Removable singularities of the solution of the harmonic map assoicated with the standard solution of the Ricci flow and removable singularities of the heat equation was studied by S.Y. Hsu in [Hs2].

Let $x_0 \in \Omega \subset \mathbb{R}^n$, $n \ge 2$, be a domain. It is well known [F] that a solution u of the harmonic equation

$$\Delta u = 0 \quad \text{in } \Omega \setminus \{x_0\}$$

has a removable singularity at x_0 if and only if

$$|u(x)| = \begin{cases} o(|x-x_0|^{2-n}) & \text{as } x \to x_0 \quad \text{if } n \ge 3 \\ o(\log|x-x_0|) & \text{as } x \to x_0 \quad \text{if } n = 2. \end{cases}$$

In this paper we will generalize the above result to the case of polyharmonic equation. We refer the reader to the book [ACL] by N. Aronszajn, T.M. Creese and L.J. Lipkin

---

1991 *Mathematics Subject Classification.* Primary 35B65 Secondary 35J30, 35J99.

Key words and phrases. removable singularity, polyharmonic equation.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

for various properties of polyharmonic functions and the papers [A], [CGS], [PV], [X], by D.H. Armitage, L.A. Cafferelli, B. Gidas and J. Spruck, J.Pipher and G.C. Verchota, and X. Xu, for some recent related results in this direction.

We now state some definitions. For any $R > 0$ and $x_0 \in \mathbb{R}^n$, let $B_R(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < R\}$ and $B_R = B_R(0)$. Let $G(x, y)$ be the Green function for the Laplacian on unit ball B_1. That is $\Delta_x G(x, y) = -\delta_y(x)$ and $G(x, y) = 0$ for all $x \in \partial B_1$, $|y| < 1$, where δ_y is the delta mass at y.

We say that a solution of of the polyharmonic equation

$$\Delta^m u = 0 \tag{1}$$

in $\Omega \setminus \{x_0\}$ has a removable singularity at x_0 if there exists a constant $\delta > 0$ such that $\overline{B_\delta(x_0)} \subset \Omega$ and a smooth solution v of (1) in $B_\delta(x_0)$ such that $u(x) = v(x)$ for all $0 < |x - x_0| < \delta$.

We choose a radially symmetric function $\phi \in C_0^\infty(\mathbb{R}^n)$, $0 \le \phi \le 1$, $\int_{\mathbb{R}^n} \phi\, dx = 1$, such that $\phi(x) = 1$ for any $|x| \le 1/2$ and $\phi(x) = 0$ for any $|x| \ge 1$. For any $\varepsilon > 0$ and function f, let $\phi_\varepsilon(x) = \varepsilon^{-n}\phi(x/\varepsilon)$ and

$$\phi_\varepsilon * f(x) = \frac{1}{\varepsilon^n}\int_{|z|\le\varepsilon} f(x - z)\phi\left(\frac{z}{\varepsilon}\right) dz.$$

We will assume $m \ge 2$ and $n \ge 2$ for the rest of the paper.

Theorem 1. *Let $x_0 \in \Omega$. A solution u of (1) in $\Omega \setminus \{x_0\}$ has a removable singularity at x_0 if and only if*

$$|\Delta^k u(x)| = \begin{cases} o(|x - x_0|^{2-n}) & \forall k = 0, 1, 2, \dots, m - 1 \quad \text{as } |x - x_0| \to 0 \quad \text{if } n \ge 3 \\ o(\log(1/|x - x_0|)) & \forall k = 0, 1, 2, \dots, m - 1 \quad \text{as } |x - x_0| \to 0 \quad \text{if } n = 2. \end{cases} \tag{2}$$

Proof. Suppose u has a removable singularity at x_0. Then there exists $\overline{B_\delta(x_0)} \subset \Omega$ and a smooth solution v of (1) in $B_\delta(x_0)$ such that $u(x) = v(x)$ for all $0 < |x - x_0| < \delta$. Hence (2) holds.

Conversely suppose (2) holds. Without loss of generality we may assume that $x_0 = 0$ and $\overline{B}_1 \subset \Omega$. For any $x \in \overline{B}_1$, let

$$v_1(x) = \int_{\partial B_1} \frac{\partial G}{\partial n_y}(x, y)\Delta^{m-1}u(y)\, d\sigma(y) \tag{3}$$

and

$$v_i(x) = \int_{B_1} G(x, y)v_{i-1}(y)\, dy + \int_{\partial B_1} \frac{\partial G}{\partial n_y}(x, y)\Delta^{m-i}u(y)\, d\sigma(y) \quad \forall i = 2, 3, \dots, m \tag{4}$$

where $\partial/\partial n_y$ is the derivative with respect to the unit inward normal at the boundary $y \in \partial B_1$ of the domain B_1. Then $v_1 \in C^\infty(B_1) \cap C(\overline{B}_1)$ and

$$\begin{cases} \Delta v_1 = 0 & \text{in } B_1 \\ v_1(x) = \Delta^{m-1}u(x) & \text{on } \partial B_1. \end{cases} \tag{5}$$

By (4) $v_2 \in C^\infty(B_1) \cap C(\overline{B}_1)$. Then by (4) and an induction argument $v_i \in C^\infty(B_1) \cap C(\overline{B})$ for all $i = 1, 2, \dots, m$. By (4),

$$v_i(x) = \Delta^{m-i}u(x) \quad \forall |x| = 1, i = 1, 2, \dots, m. \tag{6}$$

By (3),

$$\max_{|x| \le 1} |v_1| \le \max_{|x|=1} |\Delta^{m-1}u(x)|. \tag{7}$$

By (1) $\Delta^{m-1}u$ is harmonic in $B_1 \setminus \{0\}$. Hence by (2), (5), (6), and standard theory of removable singularity for harmonic functions [F],

$$\Delta^{m-1}u(x) = v_1(x) \quad \text{ in } B_1 \setminus \{0\}. \tag{8}$$

Hence $\Delta^{m-1}u$ has a removable singularity at 0 and we can extend $\Delta^{m-1}u$ to a smooth function on B_1 by letting $\Delta^{m-1}u(0) = v_1(0)$. By (4) and (6), $\forall i = 2, 3, \dots, m$,

$$\begin{aligned} \|v_i\|_{L^\infty(B_1)} \le & C_1 \|v_{i-1}\|_{L^\infty(B_1)} + \|\Delta^{m-i}u\|_{L^\infty(\partial B_1)} \\ \le & C_1(C_1\|v_{i-2}\|_{L^\infty(B_1)} + \|\Delta^{m-i+1}u\|_{L^\infty(\partial B_1)}) + \|\Delta^{m-i}u\|_{L^\infty(\partial B_1)} \\ \le & \cdots \\ \le & C_1^{i-1}\|\Delta^{m-1}u\|_{L^\infty(\partial B_1)} + C_1^{i-2}\|\Delta^{m-2}u\|_{L^\infty(\partial B_1)} + \cdots + \|\Delta^{m-i}u\|_{L^\infty(\partial B_1)} \end{aligned} \tag{9}$$

where

$$C_1 = \max_{|x| \le 1} \int_{B_1} G(x, y)\, dy = \max_{|x| \le 1} \frac{|x|^2}{2n} = \frac{1}{2n}.$$

We now claim that

$$\Delta^{m-i}u = v_i \quad \text{ in } B_1 \setminus \{0\} \tag{10}$$

for any $i = 1, 2, \dots, m$. We will prove the claim by induction. By the previous discussion the claim holds for $i = 1$. Suppose the claim holds for some $i = i_0 \in \{1, 2, \dots, m-1\}$. Then by (4),

$$\begin{aligned} & \Delta v_{i_0+1} = v_{i_0} = \Delta^{m-i_0}u \quad \text{ in } B_1 \setminus \{0\} \\ \Rightarrow \quad & \Delta(\Delta^{m-i_0-1}u - v_{i_0+1}) = 0 \quad \text{ in } B_1 \setminus \{0\}. \end{aligned} \tag{11}$$

By (2), (6), (9), (11) and standard theory of removable singularity for harmonic functions [F] we get that (10) holds for $i = i_0 + 1$. Hence by induction (10) holds for any $i = 1, 2, \dots, m$. Since $v_i \in C^\infty(B_1)$ for any $i = 1, 2, \dots, m$, by defining $\Delta^{m-i}u(0) = v_i(0)$ for all $i = 1, 2, \dots, m$, $\Delta^{m-i}u \equiv v_i \in C^\infty(B_1)$ for all $i = 1, 2, \dots, m$. Hence u has a removable singularity at $x = 0$.

Theorem 2. *Let $x_0 \in \Omega$. Suppose u is a solution of (1) in $\Omega \setminus \{x_0\}$ which satisfies*

$$|u(x)| = \begin{cases} o(|x-x_0|^{2m-n}) & \text{as } |x-x_0| \to 0 \quad \text{if } n \geq 3 \\ o(|x-x_0|^{2m-2}\log(|x-x_0|^{-1})) & \text{as } |x-x_0| \to 0 \quad \text{if } n = 2. \end{cases} \tag{12}$$

Then u has a removable singularity at x_0.

Proof. With loss of generality we may assume that $x_0 = 0$ and $\overline{B}_1 \subset \Omega$. Since u satisfies (1) in $\Omega \setminus \{0\}$, $\Delta^{m-1}u$ is harmonic in $\Omega \setminus \{0\}$. By the mean value theorem for harmonic functions (cf. Appendix C of [S]),

$$\begin{aligned} \Delta^{m-1}u(x) =& (\phi_{\frac{|x|}{2}} * \Delta^{m-1}u)(x) = ((\Delta^{m-1}\phi_{\frac{|x|}{2}}) * u)(x) \quad \forall 0 < |x| \leq 1/2 \\ \Rightarrow \quad |\Delta^{m-1}u(x)| \leq& \left(\frac{2}{|x|}\right)^n \int_{|z|\leq\frac{|x|}{2}} \left|\Delta_z^{m-1}\phi\left(\frac{z}{|x|/2}\right)\right| |u(x-z)|\,dz \\ \leq& \left(\frac{2}{|x|}\right)^{n+2(m-1)} \int_{|z|\leq\frac{|x|}{2}} \left|(\Delta^{m-1}\phi)\left(\frac{z}{|x|/2}\right)\right| |u(x-z)|\,dz \\ \leq& \frac{C}{|x|^{n+2(m-1)}} \int_{|z|\leq\frac{|x|}{2}} |u(x-z)|\,dz \quad \forall 0 < |x| \leq 1/2. \end{aligned} \tag{13}$$

By (12), for any $\varepsilon > 0$ there exists $0 < \delta_1 < 1$ such that

$$|u(x)| \leq \begin{cases} \varepsilon|x|^{2m-n} & \forall 0 < |x| \leq \delta_1 \quad \text{if } n \geq 3 \\ \varepsilon|x|^{2m-2}\log(1/|x|) & \forall 0 < |x| \leq \delta_1 \quad \text{if } n = 2 \end{cases} \tag{14}$$

Then

$$\begin{aligned} |u(x-z)| \leq& \begin{cases} \varepsilon|x-z|^{2m-n} & \forall |z| \leq \frac{|x|}{2}, 0 < |x| \leq \frac{\delta_1}{2} \quad \text{if } n \geq 3 \\ \varepsilon|x-z|^{2m-2}\log(1/|x-z|) & \forall |z| \leq \frac{|x|}{2}, 0 < |x| \leq \frac{\delta_1}{2} \quad \text{if } n = 2 \end{cases} \\ \leq& \begin{cases} C_2\varepsilon|x|^{2m-n} & \forall |z| \leq \frac{|x|}{2}, 0 < |x| \leq \delta_2 \quad \text{if } n \geq 3 \\ C_2\varepsilon|x|^{2m-2}\log(1/|x|) & \forall |z| \leq \frac{|x|}{2}, 0 < |x| \leq \delta_2 \quad \text{if } n = 2 \end{cases} \end{aligned} \tag{15}$$

for some constants $0 < \delta_2 < \delta_1/2$ and $C_2 > 0$. By (13) and (15),

$$|\Delta^{m-1}u(x)| \leq \begin{cases} C_3\varepsilon|x|^{2-n} & \forall 0 < |x| \leq \delta_2 \quad \text{if } n \geq 3 \\ C_3\varepsilon\log(1/|x|) & \forall 0 < |x| \leq \delta_2 \quad \text{if } n = 2 \end{cases} \tag{16}$$

for some constant $C_3 > 0$. Let $v_1, v_2, \dots, v_m$ be given by (3) and (4). Then by (16) and an argument similar to the proof of Theorem 1, (8) holds. Hence $\Delta^{m-1}u(x)$ has a removalbe singularity at $x = 0$. By letting $\Delta^{m-1}u(0) = v_1(0)$, $\Delta^{m-1}u(x)$ is extended to a smooth function on B_1. We now claim that both (10) and

$$|\Delta^{m-i}u(x)| = \begin{cases} o(|x|^{2-n}) & \text{as } |x| \to 0 \quad \text{if } n \geq 3 \\ o(\log(1/|x|)) & \text{as } |x| \to 0 \quad \text{if } n = 2 \end{cases} \tag{17}$$

holds for any $i = 1, 2, \ldots, m$. We will prove the claim by induction. By the previous discussion the claim holds for $i = 1$. Suppose (10) and (17) holds for some $i = i_0 \in \{1, 2, \ldots, m-1\}$. Similar to the proof of Theorem 1 (11) holds. Then by the mean value theorem for harmonic functions, $\forall 0 < |x| \leq 1/2$,

$$\begin{aligned} \Delta^{m-i_0-1}u(x) - v_{i_0+1}(x) =& \phi_{\frac{|x|}{2}} * (\Delta^{m-i_0-1}u - v_{i_0+1})(x) \\ =& \Delta^{m-i_0-1}\phi_{\frac{|x|}{2}} * u(x) - \phi_{\frac{|x|}{2}} * v_{i_0+1}(x) \\ \Rightarrow \qquad \Delta^{m-i_0-1}u(x) =& \Delta^{m-i_0-1}\phi_{\frac{|x|}{2}} * u(x) + v_{i_0+1}(x) - \phi_{\frac{|x|}{2}} * v_{i_0+1}(x). \end{aligned} \tag{18}$$

Now by (15),

$$\begin{aligned} |\Delta^{m-i_0-1}\phi_{\frac{|x|}{2}} * u(x)| \leq& \left(\frac{2}{|x|}\right)^n \int_{|z| \leq \frac{|x|}{2}} \left|\Delta_z^{m-i_0-1}\phi\left(\frac{z}{|x|/2}\right)\right| |u(x-z)|\, dz \\ \leq& \left(\frac{2}{|x|}\right)^{n+2(m-i_0-1)} \int_{|z| \leq \frac{|x|}{2}} \left|(\Delta^{m-i_0-1}\phi)\left(\frac{z}{|x|/2}\right)\right| |u(x-z)|\, dz \\ \leq& \frac{C}{|x|^{n+2(m-i_0-1)}} \int_{|z| \leq \frac{|x|}{2}} |u(x-z)|\, dz \\ \leq& \begin{cases} C\varepsilon|x|^{2(i_0+1)-n} & \forall 0 < |x| \leq \delta_2 \text{ if } n \geq 3 \\ C\varepsilon|x|^{2i_0}\log(1/|x|) & \forall 0 < |x| \leq \delta_2 \text{ if } n = 2 \end{cases} \\ \leq& \begin{cases} C'\varepsilon|x|^{2-n} & \forall 0 < |x| \leq \delta_2 \text{ if } n \geq 3 \\ C'\varepsilon\log(1/|x|) & \forall 0 < |x| \leq \delta_2 \text{ if } n = 2. \end{cases} \end{aligned} \tag{19}$$

By (9), (18) and (19), there exists a constants $C > 0$ such that

$$|\Delta^{m-i_0-1}u(x)| \leq \begin{cases} C\varepsilon|x|^{2-n} & \forall 0 < |x| \leq \delta_2 \text{ if } n \geq 3 \\ C\varepsilon\log(1/|x|) & \forall 0 < |x| \leq \delta_2 \text{ if } n = 2. \end{cases} \tag{20}$$

Hence (17) holds for $i = i_0 + 1$. By (9), (11), (20) and standard theory for removable singularity of harmonic functions, (10) holds for $i = i_0 + 1$. By induction (10) and (17) holds for all $i = 1, 2, \ldots, m$. Since $v_i \in C^\infty(B_1)$ for any $i = 1, 2, \ldots, m$, by defining $\Delta^{m-i}u(0) = v_i(0)$ for all $i = 1, 2, \ldots, m$, $\Delta^{m-i}u \equiv v_i \in C^\infty(B_1)$ for all $i = 1, 2, \ldots, m$. Hence u has a removable singularity at $x = 0$.

Remark 1. *Suppose $n \geq 3$ and either*

$$n \text{ is odd and } m \geq 1$$

or

$$n \text{ is even and } m = 1, \dots, \frac{n}{2} - 1.$$

By direct computation the function (cf. [ACL]*) $\Gamma(x) = |x|^{2m-n}$ satisfies (1) in $\mathbb{R}^n \setminus \{0\}$ and*

$$\Delta^{m-1}\Gamma(x) = A_{m,n}|x|^{2-n} \quad \forall |x| > 0 \tag{21}$$

for some constant $A_{m,n} \neq 0$. If $\Gamma(x)$ has a removable singularity at $x = 0$, then $\Gamma(x)$ can be extended to a smooth solution of (1) in a small neighborhood of 0. Hence $\Delta^{m-1}\Gamma(x) \in C^\infty$ in a small neighborhood of 0. This contradicts (21). Hence $\Gamma(x)$ has a non-removable singularity at $x = 0$. Thus Theorem 2 is sharp.

References

[A] D.H. Armitage, *On polyharmonic functions in $R^n \setminus \{0\}$*, J London Math. Soc. (2) **8** (1974), 561–569.

[ACL] N. Aronszajn, T.M. Creese and L.J. Lipkin, *Polyharmonic functions*, Clarendon Press, Oxford, 1983.

[CGS] L.A. Cafferelli, B. Gidas and J. Spruck, *Asymptotic symmetry and local behaviour of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), 271–297.

[F] G.B. Folland, *Introduction to Partial Differential Equations*, Princeton University Press and University of Tokyo Press, Princeton, New Jersey, 1976.

[H] R.S. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), 7–136, International Press, Cambridge, MA, 1995.

[Hs1] S.Y. Hsu, *Removable singularities and non-uniqueness of solutions of a singular diffusion equation*, Math. Annalen **325(4)** (2003), 665–693.

[Hs2] S.Y. Hsu, *A harmonic map flow associated with the standard solution of Ricci flow, http://arXiv.org /abs/math.DG/0702168.*

[P1] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications, http://arXiv.org /abs/math.DG/0211159.*

[P2] G. Perelman, *Ricci flow with surgery on three-manifolds, http://arXiv.org/abs/math.DG/0303109.*

[PV] J. Pipher and G.C. Verchota, *Maximum principles for the polyharmonic equation on Lipschitz domains*, Potential Analysis **4** (1995), 615–636.

[S] E.M. Stein, *Singular integral and differentiability properties of functions*, Princeton University Press, Princeton, New Jersey, 1970.

[X] X. Xu, *Uniqueness theorem for the entire positive solutions of biharmonic equations in $\mathbb{R}^n$*, Proc. Roy. Soc. Edinburgh **130A** (2000), 651–670.

E-MAIL ADDRESS:SYHSU@MATH.CCU.EDU.TW